

# Cephalopod assemblages caught by trawling along the Southern Tyrrhenian Sea (Central Mediterranean)

## Asociaciones de cefalópodos capturados por la pesca de arrastre comercial en el Tirreno meridional (Mediterráneo central)

Daniela GIORDANO\*, Teresa BOTTARI\* and Paola RINELLI\*

Recibido el 26-II-2004. Aceptado el 27-IX-2004

### ABSTRACT

The teuthofauna assemblages of the Southern Tyrrhenian Sea are identified utilising data from five trawl surveys. A total of 139 hauls were carried out in late spring or the middle of summer between 1995 and 1999, and 25 cephalopod species were found between 18 and 652 m of depth. The assemblages were analysed with the Bray-Curtis similarity index. Depth showed a positive correlation with cephalopod distribution. Four main cephalopod groups were defined: inshore (10-100 m), shelf (101-200 m), slope (201-600 m) and midslope (350-600 m). *Loligo media* was the main species for the first group, *Illex coindetii* for the second, *Todaropsis eblanae* for the third group and *Todarodes sagittatus* for the last group.

### RESUMEN

Se identifican las asociaciones de teutofauna capturadas por los arrastreros comerciales en el Tirreno meridional. Se analiza la fauna de cefalópodos recogida en 139 lances realizados en primavera o mediados del verano durante cinco campañas de pesca de arrastre realizadas entre 1995 y 1999. Se identificaron 25 especies de cefalópodos capturadas entre 18 y 652 m de profundidad. El análisis de las asociaciones faunísticas se realizó mediante el índice de similitud de Bray-Curtis. La profundidad tiene una correlación positiva con la distribución de los cefalópodos, pudiéndose definir cuatro grupos principales: costero o litoral (10-100 m), de la plataforma (101-200 m), del talud (201-600 m) y del talud medio (350-600 m). *Loligo media* fue la principal especie del primer grupo, *Illex coindetii* del segundo, *Todaropsis eblanae* del tercero, y *Todarodes sagittatus* del último grupo.

KEY WORDS: Biogeography, cephalopods, trawl fishery, Tyrrhenian Sea, faunal assemblages, Mediterranean Sea.

PALABRAS CLAVE: Biogeografía, cefalópodos, pesquería de arrastre, asociación faunística, Mar Tirreno, Mediterráneo.

### INTRODUCTION

The geographic and bathymetric distributions of demersal cephalopods have been studied in detail in different Mediterranean areas: Western Mediterranean Sea (SARTOR, BELCARI, CARBONELL, GONZÁLEZ,

QUETGLAS AND SÁNCHEZ, 1998; SORIANO, SÁNCHEZ LIZASO AND GUERRA, 2003; VILLANUEVA, 1995); the coasts of Libya and Tunisia (BONNET, 1973); Marmara Sea (KATAGAN, SALMAN AND BENLİ, 1993;

\* Istituto per l'Ambiente Marino Costiero IAMC-CNR, Messina Section, Spianata San Raineri 86, 98122 Messina, Italy. Corresponding author: Daniela Giordano. E-mail: daniela.giordano@iamc.cnr.it

UNSAL, UNSAL, ERK AND KABASAKAL, 1999); Eastern Mediterranean Sea (RUBY AND KNUDSEN, 1972; KNUDSEN, 1981; SALMAN, KATAGAN AND BOLETZKY, 1999); Ligurian Sea (RELINI, DE ROSSI, PIANO AND ZAMBONI, 2002; BERTULETTI AND ORSI RELINI, 1986; ORSI RELINI AND BERTULETTI, 1989; WURTZ, 1979; ORSI RELINI, 1995); Sardinian Sea (BONNET, 1965; CUCCU, ADDIS, DAMELE AND MANFRIN PICCINETTI, 2003); Tyrrhenian Sea (BERDAR, POTOSCHI, CAVALLARO, CAVALIERE AND LI GRECI, 1983; BERDAR AND CAVALLARO, 1975; LUMARE 1968, 1970; BELCARI, BIAGI, BIAGI, DE RANIERI, MORI AND PELLEGRINI, 1986; MANNINI AND VOLPI, 1989; BELCARI AND SARTOR, 1993; BELCARI, SARTOR AND DE RANIERI, 1998; WURTZ, MATRICARDI AND BELCARI, 1992; SPEDICATO, MINETTI, REBORA, MATRICARDI AND WURTZ, 1990; GIORDANO AND PERDICHIZZI, 1998; BELLO AND ARCULEO, 1994; GIORDANO AND CARBONARA, 1999); Sicilian Channel (JEREV AND RAGONESE, 1991, 1994; RAGONESE, JEREV AND DI STEFANO, 1992); Adriatic Sea (BELLO AND MOTOLESE, 1983; GUESCINI AND PICCINETTI MANFRIN, 1986; GAMULIN-BRIDA AND ILIJANIC, 1972; CASALI, MANFRIN PICCINETTI AND SORO, 1998; BELLO, 1990; PASTORELLI, VACCARELLA AND DE ZIO, 1995); Ionian Sea (PANETTA, D'ONGHIA, TURSI AND CECERE, 1986; TURSI AND D'ONGHIA, 1992; D'ONGHIA, MATARRESE, TURSI, MAIORANO AND PANETTA, 1995); Catalan Sea (MANGOLD-WIRZ, 1963; SÁNCHEZ, BELCARI AND SARTOR, 1998); and Aegean Sea (D'ONGHIA ET AL., 1995; SALMAN, KATAGAN AND BENLI, 2002).

In particular, assemblages of cephalopod communities have been studied in different areas of this basin (UNGARO, MARANO, MARSAN, MARTINO, MARZANO, STRIPPOLI AND VLORA, 1999; TSERPES, PERISTERAKI, POTAMIAS AND TSIMENIDES, 1999; GONZÁLEZ AND SÁNCHEZ, 2002; QUETGLAS, CARBONELL AND SÁNCHEZ, 2000). Information about spatio-temporal distributions of these species and their linkage to environmental factors is necessary for the study of many commercial species. The studies focused on demersal assemblages and in particular on their persistence, such as the capability to maintain

the same specific composition during a lag time are useful even for management purposes (BIAGI, SARTOR, ARDIZZONE, BELCARI, BELLUSCIO AND SERENA, 2002).

In this respect, this work analyses the structure of cephalopod assemblages, providing information concerning their distribution for all four groups defined: inshore, shelf, slope and mid-slope.

## MATERIAL AND METHODS

The study area is situated in the Southern part of the Tyrrhenian Sea (central Mediterranean) between Suvero Cape (Calabria) and San Vito Cape (Sicily). The data here reported come from five trawl surveys carried out from 1995 to 1999, during the MEDITS project (International bottom trawl survey in the Mediterranean Sea). The experimental surveys were always conducted during late spring or the middle of summer. Hauls were carried out from one hour before dawn until one hour after sunset. A fishing vessel equipped with an experimental trawl net with 20 mm stretched mesh size in the cod-end, and 2-2.5 m of vertical opening (FIORENTINI, COSIMI, SALA, LEONORI AND PALOMBO, 1999) was used. A total of 139 hauls were carried out, randomly allocated into five bathymetrical strata: "A": 10-50 m; "B": 51-100 m; "C": 101-200 m; "D": 201-500 m; "E": 501-800 m (BERTRAND, GIL DE SOLA, PAPACONSTANTINO, RELINI AND SOUPLET, 2002).

The duration of each haul was 30 minutes at depths less than 200 m and one hour at greater depths. All the data were standardised to one hour haul duration. All cephalopods were identified and counted on board. A multivariate approach, on the basis of the calculation of a triangular similarity matrix (group-average linkage), by depth, year and abundance (number of specimens/hour), according to BRAY AND CURTIS (1957), with relative dendrogram and non metric plan Multi Dimensional Scaling (MDS) was elaborated (CLARKE




Figure 1. Dendrogram showing similarities between hauls for the 1996 surveys. Mean depth of each haul is presented.

Figura 1. Dendrogramma que muestra las similitudes entre los arrastres de 1996. Se indica la profundidad media de cada arrastre.

AND WARWICK, 1994). Because the annual variations of cephalopod distribution were not significant, the five surveys were pooled together to calculate the total cluster. Hauls in which only one species was caught and the species with abundance values lower than 3 were not considered for the analysis. To establish which taxa contributed most to the separation of one group from another, the SIMPER routine was used (CLARKE, 1993). Abundance values were also analysed by means of univariate indices, in relation to the four assemblages evidenced by cluster analysis: total number of taxa (S), total number of individuals (N), richness of Margalef (d), Shannon-Wiener diversity (H') and Pielou's evenness (J) indices. The cumulative abundances were illustrated in order of dominance as K-dominance curves.

## RESULTS

Twenty-five species of cephalopods were collected belonging to three orders and eight families. In Table I the systematic list of species, drawn up according

to BELLO (1986), within the bathymetrical range of each species, is reported. In particular, *Onychoteuthis banksii*, *Ancistroteuthis lichtensteinii* and *Abralia veranyi* were caught only once during the period studied.

The cephalopods collected were distributed between 18 and 652 m of depth. The more coastal species was represented by *Sepia officinalis*, confined within the depth of 68 m. Fifty percent of the species showed a wide bathymetrical range that included the shelf and the beginning of the slope. The widest distribution was shown by *Eledone cirrhosa* (72-584 m), *Scaeurgus unicirrhus* (38-549 m), *Todaropsis eblanae* (61-613 m) and *Pteroctopus tetricirrhus* (118-633 m). *Alloteuthis subulata*, *Octopus vulgaris* and *Eledone moschata* were found only on the continental shelf. On the contrary, *Octopus salutii* and *Rossia macrosoma* were caught at over 200 m of depth. *Histioteuthis bonnellii* was found exclusively in the deepest stratum.

The dendrogram of all surveys (Fig. 1) showed three principal clusters: I, inshore (<100 m); II, shelf (80 - 200 m) and III, slope (200 - 600 m). The coastal group was characterized by *Loligo media*

Table I. Cephalopod species captured from 1995 to 1999 off the Southern Tyrrhenian Sea, showing their depth range and mean depth of occurrence with the associated standard deviation (SD).

Tabla I. Especies de cefalópodos capturadas desde 1995 a 1999 en el mar Tirreno meridional, indicando su rango de distribución batimétrica, profundidad media y la desviación estándar asociada (SD).

	Min	Max	Media	SD
Class CEPHALOPODA Cuvier, 1798				
Subclass COLEOIDEA Bather, 1888				
Order SEPIOIDEA Naef, 1916				
Family Sepiidae Leach, 1817				
Genus <i>Sepia</i> Linnaeus, 1758				
<i>Sepia officinalis</i> Linnaeus, 1758	40	68	49.5	12.66
<i>Sepia orbigniana</i> Féruccac in Orbigny, 1826	61	432	185.5	96.87
<i>Sepia elegans</i> Blainville, 1827	40	345	137.3	74.02
Family Sepiolidae Leach, 1817				
Subfamily Sepiolinae Leach, 1817				
Genus <i>Sepiola</i> Leach, 1817				
<i>Sepiola randeletii</i> Leach, 1817	110	395	276	115
Genus <i>Rondeletiola</i> Naef, 1921				
<i>Rondeletiola minor</i> (Naef, 1912)	117	297	207	127.28
Subfamily Rossiinae Appelof, 1898				
Genus <i>Rossia</i> Owen in Ross, 1835				
<i>Rossia macrosoma</i> (Delle Chiaie, 1830)	277	596	484.5	115.56
Genus <i>Neorossia</i> Boletzky, 1971				
<i>Neorossia caroli</i> (Joubin, 1902)	123	608	309	195.9
Order TEUTHOIDEA Naef, 1916				
Suborder MYOPSIDA Orbigny in Féruccac & Orbigny, 1840				
Family Loliginidae Lesueur, 1821				
Genus <i>Loligo</i> Lamarck, 1798				
<i>Loligo vulgaris</i> Lamarck, 1798	18	294	75.72	72
<i>Loligo forbesi</i> Steenstrup, 1856	72	505	134.2	127.3
<i>Loligo media</i> (Linnaeus, 1758)	18	383	99.1	75.43
Genus <i>Alloteuthis</i> Wukler, 1920				
<i>Alloteuthis subulata</i> (Lamarck, 1798)	76	143	109.5	47.4
Suborder OEGOPSIDA Orbigny, 1845				
Family Enoploteuthidae Pfeffer, 1900				
Subfamily Enoploteuthinae Pfeffer, 1900				
Genus <i>Abralia</i> Gray, 1849				
<i>Abralia veranyi</i> (Ruppell, 1844)	316	316	-	-
Family Onychoteuthidae Gray, 1847				
Genus <i>Onychoteuthis</i> Lichtenstein, 1818				
<i>Onychoteuthis banksii</i> (Leach, 1817)	358	358	-	-
Genus <i>Ancistroteuthis</i> Gray, 1849				
<i>Ancistroteuthis lichensteinii</i> (Féruccac, 1839)	562	562	-	-
Family Histioteuthidae Verrill, 1881				
Genus <i>Histioteuthis</i> Orbigny, 1840				
<i>Histioteuthis bonnellii</i> (Féruccac, 1834)	531	645	565.5	53.33
<i>Histioteuthis reversa</i> (Verrill, 1880)	383	652	565.3	89.4
Family Ommastrephidae Steenstrup, 1857				
Subfamily Illicinae Posselt, 1890				
Genus <i>Illex</i> Steenstrup, 1880				
<i>Illex coindetii</i> (Verany, 1839)	60	593	213.7	128.7

Table I. Continuación.  
Tabla I. Continuation.

		Min	Max	Media	SD
Genus <i>Todaropsis</i> Girard, 1890					
<i>Todaropsis eblanae</i> (Boll, 1841)		61	613	323.1	152.2
Subfamily Todardinae Adam, 1960					
Genus <i>Todarodes</i> Steenstrup, 1880					
<i>Todarodes sagittatus</i> (Lamarck, 1798)		114	593	438.5	154.69
Order OCTOPODA Leach, 1818					
Suborder INCIRRATA Grimpe, 1916					
Family Octopodidae Orbigny in Féussac & Orbigny, 1840					
Subfamily Octopodinae Orbigny in Féussac & Orbigny, 1840					
Genus <i>Octopus</i> Cuvier, 1798					
<i>Octopus vulgaris</i> Cuvier, 1798		27	114	62.86	25.85
<i>Octopus salutii</i> Vérany, 1836		295	432	374	70.87
Genus <i>Scaeurgus</i> Troschel, 1857					
<i>Scaeurgus unicirrhus</i> (Delle Chiaje in Féussac & Orbigny, 1840)		38	549	174.1	110.8
Genus <i>Pteroptopus</i> P. Fischer, 1882					
<i>Pteroptopus tetricirrus</i> (Delle Chiaje, 1830)		118	633	353.8	147.83
Subfamily Eledoninae Grimpe, 1921					
Genus <i>Eledone</i> Leach, 1817					
<i>Eledone moschata</i> (Lamarck, 1798)		18	143	115.4	92.82
<i>Eledone cirrhosa</i> (Lamarck, 1798)		72	584	200.4	119.87

and *Loligo vulgaris*. In the shelf group (Table IIa) *Illex coindetii* and *Loligo media* prevailed, followed by *Sepia orbignyana*, *Sepia elegans* and *Scaeurgus unicirrhus*. The slope group (Table IIb) was characterized by *Todaropsis eblanae*. The Multi-dimensional Scaling analysis (30%: Stress 0.15) also showed four groups, well discriminated by depth (Fig. 2). There is also a midslope group (Fig. 3), limited to a few hauls, in which *Todarodes sagittatus* was the prevalent species.

Analysing the univariate indices (Table III), elaborated for the four groups, the highest biodiversity was observed in the shelf. In this macrostratum, the highest values of Margalef and Shannon-Wiener indices were recorded as well as the highest number of species and individuals. The highest values of H' found at this bathymetrical level can be explained by the relative homogeneity in abundance of the most frequent species.

However, a trend was observed according to the depth as evidenced also by the cumulative abundance curve, in which the shelf group showed the highest biodiversity and the midslope the lowest. In this last group in fact both d and H' values recorded the lowest values (0.973 and 1.063, respectively).

Finally the Evenness (J') values ranged from 0.616 to 0.661. The higher value was recorded for the last cluster. Also in the second cluster this value is quite high.

## DISCUSSION AND CONCLUSIONS

The 25 cephalopod species distributed in 8 families recorded in this study account for 42.4% of species reported from the Mediterranean (MANGOLD AND BOLETZKY, 1988; GUERRA, 1992). Absent were certain species of the Sepiolidae family and pelagic species. In compari-

Table IIa. Indicator species and related data from the SIMPER analysis. Abbreviations: AA, average abundance, contribution of each species in each group; AT, average term, average Bray-Curtis contribution of each species to distinguish between groups. The ratio (AT/SD), the percentage contribution to the separation (%), and the cumulative percentage (Cum %) are shown for each group comparison.

Tabla IIa. Especies indicadoras y datos relacionados, según el análisis SIMPER. Abreviaturas: AA, abundancia promedio, contribución de cada especie en cada uno de los grupos; Average Term (AT), contribución promedio de Bray-Curtis de cada especie en la diferenciación de los grupos. Cada grupo viene definido, además, por la relación AT/SD (ratio), la contribución porcentual (%), y el porcentaje acumulativo de cada especie (Cum %).

Average dissimilarity: 84.1	inshore AA	shelf AA	AT	Ratio	%	Cum (%)
<i>I. vulgaris</i>	106.88	0.90	34.64	1.39	41.18	41.18
<i>I. coindetii</i>	0.58	26.62	0.50	0.67	13.74	54.92
<i>L. vulgaris</i>	18.88	0.02	0.47	1.03	13.22	68.13
<i>S. elegans</i>	0.05	0.41	0.28	0.71	7.70	75.83
<i>S. orbignyana</i>	0.00	8.86	5.69	0.53	6.77	82.6
<i>S. unicirrus</i>	0.00	0.29	0.21	0.6	5.45	88.05
<i>E. cirrhosa</i>	0.05	0.13	0.13	0.6	3.57	91.62
Average dissimilarity: 99	inshore AA	shelf AA	AT	Ratio	%	Cum (%)
<i>L. media</i>	106.88	0.00	47.82	1.48	48.3	48.3
<i>L. vulgaris</i>	18.88	0.00	0.80	1.07	19.25	67.56
<i>T. eblanae</i>	0.00	0.46	9.70	0.64	9.8	77.35
<i>P. tetricirrus</i>	0.00	2.74	3.81	0.67	3.85	81.2
<i>O. vulgaris</i>	0.09	0.00	0.13	0.63	3.17	84.37
<i>S. orbignyana</i>	0.00	2.89	2.68	0.32	2.71	87.08
<i>I. coindetii</i>	0.04	1.67	2.43	0.56	2.46	89.54
<i>E. moschata</i>	2.00	0.19	2.37	0.76	2.39	91.93
Average dissimilarity: 92.7	inshore AA	shelf AA	AT	Ratio	%	Cum (%)
<i>L. media</i>	21.43	0.00	19.36	1.03	20.89	20.89
<i>I. coindetii</i>	26.62	1.67	17.55	0.82	18.94	39.83
<i>S. elegans</i>	9.57	0.00	10.85	0.82	11.71	51.54
<i>S. orbignyana</i>	8.86	2.89	10.51	0.69	11.34	62.88
<i>T. eblanae</i>	1.14	10.56	9.67	0.70	10.43	73.31
<i>S. unicirrus</i>	6.54	0.85	7.77	0.69	8.38	81.69
<i>E. cirrhosa</i>	3.30	0.67	4.95	0.63	5.34	87.04
<i>P. tetricirrus</i>	0.22	2.74	3.47	0.83	3.75	90.79

son, TURSI AND D'ONGHIA (1992) caught 24 species of cephalopods in the Ionian Sea, D'ONGHIA ET AL. (1995) 29 species in the North Aegean Sea, QUETGLAS ET AL. (2000) found 30 species in the Balearic Sea, SÁNCHEZ ET AL. (1998) found 36 species in the northern Tyrrhenian Sea and 47 in the Catalan Sea and SORIANO ET AL. (2003) found 18 species in the upper continental slope of Alicante (western Mediterranean).

In previous surveys of the study area before 1995 (GIORDANO AND PERDICHIZZI, 1998), 32 species of cephalopods were identified. In general, a similar cephalopod distribution pattern was observed from 1994-1995 trawl surveys, but the sampling was carried out at a different period (in autumn) and with a different net (with 36 mm of stretched mesh). The species caught in the previous faunistic list in the same area were: *Sepiola affinis*,

Table IIb.  
Tabla IIb.

Average dissimilarity: 100	inshore AA	midslope AA	AT	Ratio	%	Cum (%)
<i>L. media</i>	106.88	0	50.79	1.52	50.79	50.79
<i>L. vulgaris</i>	18.88	0	21.3	1.08	21.3	72.09
<i>T. sagittatus</i>	0	7	9.23	0.85	9.23	81.32
<i>O. vulgaris</i>	2.5	0	3.63	0.62	3.63	84.94
<i>S. orbignyana</i>	0	3.6	3.51	0.42	3.51	88.46
<i>E. moschata</i>	2	0	2.61	0.74	2.61	91.07
Average dissimilarity: 97.18	inshore AA	midslope AA	AT	Ratio	%	Cum (%)
<i>L. media</i>	21.43	0	20.96	1.05	21.57	21.57
<i>I. coindetii</i>	26.62	0	18.87	0.81	19.42	40.98
<i>S. elegans</i>	9.57	0	11.87	0.84	12.21	53.19
<i>S. orbignyana</i>	8.86	3.6	11.6	0.72	11.93	65.13
<i>T. sagittatus</i>	0.16	7	8.66	0.99	8.91	74.04
<i>S. unicirrus</i>	6.54	0.2	8.41	0.68	8.66	82.7
<i>E. cirrhosa</i>	3.3	0	5.64	0.61	5.8	88.5
<i>E. moschata</i>	0.22	0	2.65	0.64	2.72	91.22
Average dissimilarity: 92.38	inshore AA	midslope AA	AT	Ratio	%	Cum (%)
<i>T. eblanae</i>	10.56	0	23.25	0.95	25.17	25.17
<i>T. sagittatus</i>	0.3	7	21.55	1.3	23.33	48.5
<i>P. tetricirrus</i>	2.81	0.4	11.48	0.96	12.42	60.92
<i>S. orbignyana</i>	2.22	3.6	10.36	0.57	11.21	72.14
<i>R. macrosoma</i>	0.19	1	6.78	0.6	7.34	79.48
<i>I. coindetii</i>	1.67	0	5.67	0.6	6.14	85.61
<i>S. unicirrus</i>	0.85	0.2	3.49	0.57	3.78	89.39
<i>E. cirrhosa</i>	0.67	0	2.93	0.46	3.17	92.56

*Sepiola ligulata*, *Sepiola robusta*, *Sepiella oweni*, *Sepiella neglecta* (Sepiolidae), *Chiroteuthis veranii* (Chiroteutidae) and *Bathypolypus sponsalis* (Octopodidae). On-board observation of catches indicated that the demersal fauna sampled is strongly influenced by depth. The

depth range of each species with mean depth and standard deviation are reported in Table I. *Eledone cirrhosa* (32-584 m), *Scaeurgus unicirrus* (38-549 m), *Todaropsis eblanae* (61-613 m), *Sepia elegans* (40-432 m), *Loligo media* (18-383 m) and *Illex coindetii* (60-593 m) showed

Table III. Univariate diversity indexes for each cluster. Abbreviations: S, number of species; N, number of specimens; d, richness of Margalef; J, Pielou's evenness; H', Shannon-Wiener diversity. Tabla III. Índices univariados de diversidad de cada grupo. Abreviaturas: S, número de especies; N, número de ejemplares; d, riqueza según Margalef; J, uniformidad según Pielou; H', diversidad de Shannon-Wiener.

Cluster	S	N	d	J'	H'
inshore	8	709	1.066	0.616	1.28
shelf	15	2267	1.812	0.652	1.765
slope	12	529	1.754	0.619	1.539
midslope	5	61	0.973	0.661	1.063




Figure 2. Multidimensional scaling ordination analysis for the five surveys.

Figura 2. Análisis de ordenación de escalamiento multidimensional para las cinco campañas.

a wide distribution range, extending up to the continental slope.

During the present study four main cephalopod groups were defined: inshore (10-100 m), shelf (101-200 m), slope (201-600 m) and midslope (350-600 m). *Loligo media* was the main species for the first group, *Illex coindetii* for the second, *Todaropsis eblanae* for the

third group and *Todarodes sagittatus* for the last group.

## ACKNOWLEDGEMENTS

We are grateful to Dr. Patrizia Jereb for helpful comments on the manuscript.

## BIBLIOGRAPHY

- BELCARI, P., BIAGI, F., BIAGI, V., DE RANIERI, S., MORI, M. AND PELLEGRINI, D., 1986. Observations about Cephalopods distribution in the north Tyrrhenian sea. *Rapport de la Commission Internationale pour l'exploration scientifique de la Mer Méditerranée*, 30 (2): 246.
- BELCARI, P. AND SARTOR, P., 1993. Bottom trawling teuthofauna of the North Tyrrhenian sea. *Scientia Marina*, 57 (3): 145-152.
- BELCARI, P., SARTOR, P. AND DE RANIERI, S., 1998. I cefalopodi nello sbarcato commerciale con reti a strascico nel Mar Tirreno settentrionale. *Biologia Marina Mediterranea*, 5 (2): 318-325.
- BELLO, G., 1986. Catalogo dei molluschi cefalopodi viventi nel Mediterraneo. *Bollettino Malacologico*, 22: 197-214.
- BELLO, G., 1990. The Cephalopods fauna of the Adriatic. *Acta Adriatica*, 31: 275-291.
- BELLO, G. AND ARCULEO, M., 1994. I cefalopodi dei fondi strascicabili del Golfo di Castellammare. *Bollettino Malacologico*, 30 (5-9): 173-181.
- BELLO, G. AND MOTOLESE, G. 1983. Sepiolidi from the Adriatic Sea (Mollusca, Cephalopoda). *Rapport de la Commission Internationale pour l'exploration scientifique de la Mer Méditerranée*, 28 (5): 281-282.
- BERDAR, A., POTOSCHI, A., CAVALLARO, G., CAVALIERE, A. AND LI GRECI, F., 1983. Su alcuni cefalopodi spiaggiati e pescati nello Stretto di Messina. *Memorie di Biologia Marina ed Oceanica*, 13 (2): 115-127.
- BERDAR, A. AND CAVALLARO, G., 1975. Ulteriore contributo alla conoscenza dei Cefalopodi spiaggiati lungo la costa siciliana dello Stretto di Messina. *Memorie di Biologia Marina ed Oceanica*, 5 (5): 121-138.
- BERTRAND, J. A., GIL DE SOLA, L., PAPACONSTANTINOU, C., RELINI, G. AND SOUPLET, A., 2002. An international bottom trawl in the Mediterranean: the MEDITS programme. *1995 ICES Annual Science Conference*, CM 1997/y:03: 16 pp.

- BERTULETTI, M. AND ORSI RELINI, L., 1986. Notes on the presence and distribution of the Sepiolinae in the Ligurian Sea. *Rapport Commission Internationale pour l'exploration Scientifique de la Mer Méditerranée*, 30 (2): 247.
- BIAGI, F., SARTOR, P., ARDIZZONE, G., BELCARO, P., BELLUSCIO, A. AND SERENA, F., 2002. Analysis of demersal assemblages off the Tuscany and Latium coasts (north-western Mediterranean). *Scientia Marina*, 66 (suppl. 2): 233-242.
- BONNET, M., 1973. Les céphalopodes capturés par la Thalassa en Novembre 1969 au large de la Libye et de la côte orientale tunisienne. *Revue des travaux de l'Institut des Pêches Maritimes*, 37 (2): 253-256.
- BONNET, M., 1965. Remarques sur l'écologie des Céphalopodes des côtes de Sardegna et de Corse capturés par la "Thalassa" en novembre et décembre 1963. *Rapport de la Commission Internationale pour l'exploration scientifique de la Mer Méditerranée*, 18: 234-240.
- BRAY, J. R. AND CURTIS, J. T., 1957. An ordination of the upland forest communities of the Southern Wisconsin. *Ecological Monographs*, 27: 325-349.
- CASALI, P., MANFRIN PICCINETTI, G. AND SORO, S., 1998. Distribuzione di cefalopodi in alto e medio Adriatico. *Biologia Marina Mediterranea*, 5 (2) : 307-317.
- CLARKE, K. R., 1993. Non parametric multivariate analyse of changes in community structure. *Australian Journal of Ecology*, 18: 117-143.
- CLARKE, K. R. AND WARWICK, R. M., 1994. *Change in marine communities: an approach to statistical analysis and interpretation*. Plymouth Marine Laboratory, Plymouth, 144 pp.
- CUCCU, D., ADDIS, P., DAMELE, F. AND MANFRIN PICCINETTI, G., 2003. Primo censimento della teutofauna dei mari circostanti la Sardegna. *Biologia Marina Mediterranea*, 10 (2):795-798.
- D'ONGHIA, G., MATARRESE, A., TURSI, A., MAIORANO P. AND PANETTA, P., 1995. Osservazioni sulla teutofauna epi e mesobatiale nel Mediterraneo Orientale (Mar Ionio e Mar Egeo). *Biologia Marina Mediterranea*, 2 (2): 199-204.
- FIORENTINI, L., COSIMI, G., SALA, A., LEONORI, I. AND PALOMBO, V., 1999. Efficiency of the bottom trawl used for Mediterranean international trawl survey (MEDITS). *Aquatic Living Resources*, 12 (3): 187-205.
- GAMULIN-BRIDA, H. AND ILIJANIC, V., 1972. Contribution à la connaissance des Céphalopodes de l'Adriatique. *Acta Adriatica*, 14 (6) : 12 pp.
- GIORDANO, D. AND CARBONARA, P., 1999. Nota sulla distribuzione di molluschi cefalopodi nel Tirreno centro-meridionale. *Biologia Marina Mediterranea*, 6 (1): 573-575.
- GIORDANO, D. AND PERDICHIZZI, F., 1998. Osservazioni sulla teutofauna nel Tirreno meridionale da Capo Suvero (Calabria) a Capo S. Vito (Sicilia). *Biologia Marina Mediterranea*, 5 (1): 807-810.
- GONZÁLEZ, M. AND SÁNCHEZ, P., 2002. Cephalopod assemblages caught by trawling along the Iberian Peninsula Mediterranean coast. *Scientia Marina*, 66 (Suppl. 2): 199-208.
- GUERRA, A., 1992. Mollusca Cephalopoda. En: *Fauna Ibérica*, vol. 1. Ramos, M. A. et al. (Ed.). Museo Nacional de Ciencias Naturales, CSIC, Madrid, 327 pp.
- GUESCINI, A. AND PICCINETTI MANFRIN, G., 1986. Distribuzione di Sepiolidi nell'Adriatico centro-settentrionale. *Nova Thalassia*, 8, Suppl. 3, pp. 513-518.
- JEREB, P. AND RAGONESE, S., 1991. On the association of the squid *Illex coindetii* (Mollusca: Cephalopoda) with the target species trawled in the Sicilian Channel. *Note Tecniche e Reprints - Istituto di Tecnologia della Pesca e del Pescato*, 32: 3 p.
- JEREB, P. AND RAGONESE, S., 1994. The Mediterranean teuthofauna: Towards a biogeographical coverage by regional census. II. Strait of Sicily. *Bollettino Malacologico*, 30 (5-9): 161-172.
- KATAGAN, T., SALMAN, A. AND BENLI, H. A., 1993. The cephalopod fauna of the Sea of Marmara. *Israel Journal of Zoology*, 39: 255-261.
- KNUDSEN, J., 1981. Three Sepiolidae to the eastern Mediterranean (Mollusca: Cephalopoda). *Argamon, Israel Journal of Malacology*, 7: 45-50.
- LUMARE, F., 1968. Ricerche sui Cefalopodi dell'alto Tirreno. *Commissione Studi Oceanografici e di Limnologia, Programma di ricerca sulle risorse Marine e del fondo Marino*, C. N. R. B., 24: 5-29.
- LUMARE, F., 1970. Nota sulla distribuzione di alcuni cefalopodi del mar Tirreno. *Bollettino Pesca Piscicoltura e Idrobiologia*, 25 (2): 313-344.
- MANGOLD, K. AND BOLETSKY, S., 1988. Mediterranean Cephalopod Fauna. In: M. R. Clarke and E. R. Trueman (eds.), *Ethology and Neontology of Cephalopods, The Mollusca*, 12: 315-330. Academic Press, San Diego.
- MANGOLD-WIRZ, K., 1963. Biologie des Céphalopodes benthiques et nectoniques de la Mer Catalane. *Vie et Milieu*, 13: 1-285.
- MANNINI, P. AND VOLPI, C., 1989. Nota sulla presenza e distribuzione di alcuni cefalopodi del Tirreno settentrionale. *Oebalia*, 25-2, N. S.: 693-701.
- ORSI RELINI, L., 1995. Notes on midwater collections of *Heteroteuthis dispar* (Cephalopoda, Sepiolidae). *Bulletin de l'Institut océanographique Monaco*, n. spécial 16 : 63-72.

- ORSI RELINI, L. AND BERTULETTI, M., 1989. Sepiolines (Mollusca, Cephalopoda) from the Ligurian Sea. *Vie et Milieu*, 39 (3/4):183-190.
- PANETTA, P., D'ONGHIA, G., TURSI, A. AND CECERE, E., 1986. Il ruolo dei Molluschi Cefalopodi nella valutazione delle risorse marine dello Jonio (campagne di pesca 1985). *Nova Thalassia*, 8, Suppl. 3:523-527.
- PASTORELLI, A. M., VACCARELLA, R. AND DE ZIO, V., 1995. Distribuzione dei Cefalopodi commerciali del basso Adriatico. *Biologia Marina Mediterranea*, 2 (2): 501-502.
- QUETGLAS, A., CARBONELL, A. AND SÁNCHEZ, P., 2000. Demersal continental shelf and upper slope cephalopods assemblages from the Balearic Sea (North-Western Mediterranean). Biological aspects of some deep-sea species. *Estuarine, Coastal and Shelf Science*, 50: 739-749.
- RAGONESE, S., JEREB, P. AND DI STEFANO, L., 1992. Occurrence of the unicorn octopus *Scaeurgus unicirrhus* in the Sicilian Channel. I. Spatial distribution and abundance. II: Bathymetric distribution. *Rapport Commission Internationale pour l'exploration Scientifique de la Mer Méditerranée*, 33: 306-307.
- RELINI, G., DE ROSSI, C., PIANO, T. AND ZAMBONI A., 2002. Osservazioni sui Cefalopodi dei fondi strascicabili liguri. *Biologia Marina Mediterranea*, 9 (1): 792-795.
- RELINI, G. AND ORSI RELINI, L., 1984. The role of cephalopods in the inshore trawl fishing of the Ligurian Sea. *Oebalia*, 10: 35-58.
- RESTUCCIA, V. AND RAGONESE, S., 1986. Cefalopodi dei fondali strascicabili della Sicilia nord-orientale: nota preliminare. *Nova Thalassia*, 8, Suppl. 3:543-549.
- RUBY, G. AND KNUDSEN, J., 1972. Cephalopoda from the Eastern Mediterranean. *Israel Journal of Zoology*, 21: 83-97.
- SALMAN, A., KATAGAN, T. AND BENLI, H. A., 2002. Cephalopod Fauna of the Eastern Mediterranean. *Turkish Journal of Zoology*, 26: 47-52.
- SALMAN, A., KATAGAN, T. AND BOLETZKY, S. V., 1999. New cephalopod molluscs in the Eastern Mediterranean: previously unnoted species or Lessepsian migrants? *Vie et Milieu*, 49 (1): 11-17.
- SÁNCHEZ, P., BELCARI, P. AND SARTOR, P., 1998. Composition and spatial distribution of cephalopods in two north-western Mediterranean areas. *South African Journal of Marine Science*, 20: 17-24.
- SARTOR, P., BELCARI, P., CARBONELL, A., GONZÁLEZ, M., QUETGLAS, A. AND SÁNCHEZ, P., 1998. The importance of cephalopods to trawl fisheries in the western Mediterranean. *South African Journal of Marine Science*, 20: 67-72.
- SORIANO, S., SÁNCHEZ LIZASO, J. L. AND GUERRA, A., 2003. Cephalopod species collected in the upper continental slope off Alicante (Western Mediterranean). *Iberus*, 21: 57-66.
- SPEDICATO, M. T., MINETTI, D., REBORA, F., MATRICARDI, G. AND WURTZ, M., 1990. Distribuzione di Cefalopodi Decapodi nel medio-basso Tirreno. *Oebalia*, Suppl. 26 (2): 761-766.
- TSERPES, G., PERISTERAKI, P., POTAMIAS, G. AND TSIMENIDES, N., 1999. Species distribution in the southern Aegean sea based on bottom-trawl surveys. *Aquatic Living Resources*, 12 (3): 167-175.
- TURSI, A. AND D'ONGHIA, G., 1992. Cephalopods in the Ionian Sea (Mediterranean sea). *Oebalia*, 18: 25-43.
- UNGARO, N., MARANO, C. A., MARSAN, R., MARTINO, M., MARZANO, M. C., STRIPPOLI, G. AND VLORA, A., 1999. Analysis of demersal species assemblages from trawl surveys in the South Adriatic sea. *Aquatic Living Resources*, 12 (3): 177-185.
- UNSAK, I., UNSAL, N., ERK, M. H. AND KABA-SAKAL, H., 1999. Demersal cephalopods from the Sea of Marmara with remarks on some ecological characteristics. *Acta Adriatica*, 40: 105-104.
- VILLANUEVA, R., 1995. Distribution and abundance of bathyal sepiolids (Mollusca: Cephalopoda) in the northwestern Mediterranean. *Bulletin de l'Institut océanographique, Monaco*, n° special 16: 19-26.
- VOLPI, C., AUTERI, R., BORRI, M. AND MANNINI, P., 1990. Distribution and reproduction of *Sepia elegans* in the North Tyrrhenian Sea. *Rapport Commission Internationale pour l'exploration Scientifique de la Mer Méditerranée*, 32 (1): 244.
- WURTZ, M., 1979. I Cefalopodi raccolti in Mar Ligure durante la campagna di pesca batiale 1977/1978. *Atti della Società Toscana di Scienze Naturali, memorie*, ser. B, 86: 374-377.
- WURTZ, M., MATRICARDI, G. AND BELCARI, P., 1992. Distribution and abundance of the Octopus *Eledone cirrhosa* in the Tyrrhenian sea, central Mediterranean. *Fishery Research*, 13 (1): 53-66.